



## PREVALENCE OF INFECTIOUS DISEASES AND DISORDERS IN RELATION TO HEAD AND NECK REGION. PERTAINING TO AN ORIGINAL RESEARCH

**Dr. Susmita Choudhary<sup>1</sup>, Dr. Shivangi Verma<sup>2</sup>, Dr Anindita Talukdar<sup>3</sup>, Dr. Karthik Shunmugavelu<sup>4</sup>, Dr. Evangeline Cynthia Dhinakaran<sup>5</sup>,**

**Dr. Datta Sai Kiran<sup>6</sup>**

1. Associate Professor, Department of Orthodontics and Dentofacial Orthopaedics  
Narsinhbhai Patel Dental College and Hospital, Visnagar, Gujarat 384315, India.

2 BDS, MDS Senior lecturer Department of Paediatric Dentistry RKDF dental college  
Bhopal Madhya Pradesh, India

3. Assistant Professor, Department of Pedodontics and Preventive Dentistry Regional  
Dental College, Guwahati, Assam. India.

4. PSP Medical College Hospital and Research Institute. Tambaram Kanchipuram main  
road Oragadam Panruti Kanchipuram district Tamilnadu 631604.

5. MBBS MD PATHOLOGY. Senior Resident Department of Pathology. Sree Balaji  
Medical College and Hospital, Chrompet, Chennai-600044, Tamilnadu,

India.

6. Sr.lecturer, G Pulla Reddy Dental College and Hospital

Received: 18/12/2024  
Accepted: 21/01/2025

EMAIL: [drkarthiks1981@gmail.com](mailto:drkarthiks1981@gmail.com)



## ABSTRACT

Assessment of oral health includes careful examination of hard and soft tissues. The Global Burden of Diseases, 2015 indicates that dental caries and periodontal diseases are the most common oral disorders followed by other oral lesions. The aim of the study is to identify the pattern and expression of oral mucosal lesions using WHO guidelines and Oral Hygiene Index – S respectively in paediatric population. 100 patients 3-16 years of age from department of paediatrics in a multispeciality hospital, Chennai were examined. Majority of them had oral lesions (68%) with traumatic ulceration being the most common.

**KEYWORDS:** oral; paediatric; dental; pathology; lesions.

## PREVALENCIA DE ENFERMEDADES Y TRASTORNOS INFECCIOSOS EN RELACIÓN CON LA REGIÓN DE LA CABEZA Y EL CUELLO. UNA INVESTIGACIÓN ORIGINAL

## RESUMEN

La evaluación de la salud bucal incluye un examen cuidadoso de los tejidos duros y blandos. La Carga Global de Enfermedades de 2015 indica que la caries dental y las enfermedades periodontales son los trastornos bucales más comunes, seguidos de otras



lesiones bucales. El objetivo del estudio es identificar el patrón y la expresión de las lesiones de la mucosa oral utilizando las directrices de la OMS y el Índice de Higiene Bucal – S respectivamente en población pediátrica. Se examinaron 100 pacientes de 3 a 16 años de edad del departamento de pediatría de un hospital multiespecializado de Chennai. La mayoría presentaba lesiones orales (68%), siendo la ulceración traumática la más común.

**PALABRAS CLAVE:** oral; pediátrico; dental; patología; lesiones.

## INTRODUCTION

Oral and dental health does not comprise only dental caries and periodontal diseases<sup>1</sup>. Oral mucosal lesions (OML) present as alterations in the soft tissue of the oral cavity associated with etiopathogenesis, clinical features and variable diagnostic and prognostic characteristics<sup>2</sup>. Oral mucosal lesions can be benign or potentially malignant requiring no treatment to extensive invasive treatment<sup>3</sup>. The recognition of

these lesions involves thorough history taking and intraoral examination<sup>4</sup>. The assessment of soft tissues in paediatric patients involves knowledge of normal size; shape, color, and texture of the structures that comprise it<sup>2</sup>. Epidemiological studies have shown significant variation among different geographical locations<sup>5</sup>. Literature evidence of oral lesions among the pediatric population is scarce. The presentation of lesions in oral mucosa in



children can differ from adults in respect to color, size alterations, aetiology, clinical characteristics, prognosis, and treatment protocol. This it is importance to assess the prevalence of the oral lesions in pediatric population for appropriate management. The main objective of this study was to identify the pattern and expression of oral mucosal lesions using WHO guidelines and Oral Hygiene Index – S respectively in pediatric population.

## **MATERIALS AND METHODS**

The study sample includes 100 patients 3-16 years of age from department of pediatrics in a multispeciality hospital, Chennai. A thorough history taking was done. Demographics, chief complaint, history of presenting illness, past medical

history, extra oral and intra oral examination were assessed. Intraoral examination involved assessment of hard tissue, soft tissue and radiographic examination. The lesions were recorded in a structured format involving site, size, shape, color, consistency and extension. This study was done over a period of one year between April 2023 to April 2024. The inclusion criteria included outpatients and inpatients of 3 - 16 years of age. The exclusion criteria were patients who were not co-operative and patients who are unable to participate due to systemic illness. The results were collected and analysed with chi-squared test and ANOVA using Statistical Package for Social Sciences (SPSS version 21).



## RESULTS

### *Demographics*

The gender distribution was 52% male and 48% female. When the relationship between age and BMI were assessed majority of them (54%) were normal, 26% were underweight, 11% were overweight and 9% of the children were obese. Majority of the children examined had good OHI score.

### *Prevalence of lesion*

Majority of the lesions present between 3-4 years of age ( $p=.001$ ) and the least in the age group of 15-16 years of age. The lesions were present on 68% of the population. The most common lesion observed was traumatic ulceration, followed by dentoalveolar abscess, gingivitis, geographic tongue, while the

other lesions comprising of white lesion, ankyloglossia, eruption cyst and aphthous ulcer encased 1% of the oral lesions present in the patients examined.

## DISCUSSION

In this study, oral screening was done for 100 children. 68% of the participants had lesions whereas 32% were devoid of lesions. Male participants recorded higher oral mucosal lesions. 44% of the lesions found in age group of 3-4 years. A majority of the patients examined had oral lesions, of which traumatic ulcers were the most predominant.

Shulman et al examined the prevalence of oral mucosal lesions in children and youths in the USA and found a majority of them had lesions present with lip being



the most common site and lip bite/cheek bite, the most common lesion. Males had more lesions present compared to females<sup>6</sup>.

Hussein et al assessed the prevalence of oral lesions among the Jordanian children and found that 47.4% had oral lesions and that there was no significant difference among the gender. However, he deduced that the prevalence increased with age<sup>7</sup>.

Ambika et al examined oral lesions in children attending an Indian public school and observed its presence in 64.11% of the sample population. The most common lesions evident were gingivitis, gingival abscess and traumatic ulcers<sup>8</sup>.

#### *Traumatic ulcer*

Ulcerations can cause defects in the epithelium, connective tissue or both.

Traumatic ulceration is a common oral mucosal lesion due to habits, malocclusion, sharp tooth, mechanical/chemical/thermal injury and vitamin deficiency. With their location being buccal mucosa, labial mucosa. They can be solitary or multiple. They can persist for a few days or a few weeks but become painless three days after elimination of the injury and heal in 10 days<sup>9</sup>.

#### *Geographic tongue*

It is a benign recurrent condition affecting the tongue with loss of epithelium. The aetiology is unknown with burning sensation, pain or it can be asymptomatic. It persists for a few days to few weeks and can reappear at a different location<sup>10</sup>.



### *Gingivitis*

Gingival inflammation in children can progress to cause destruction of the periodontium of the adult. The wide contact points between the interdental region in children, increases the susceptibility to bacterial growth and provide for a wider area of destruction<sup>11</sup>. Abrams et al conducted a study on prevalence of gingivitis among well and malnourished children and found that no significant difference in the Plaque Index (PII) and the Modified Gingival Index (MGI) among well-nourished and malnourished groups and between males and females. But there was less plaque and gingivitis among well nourished children when examined with age percentiles<sup>12</sup>.

### *Dentoalveolar abscess*

Dentoalveolar abscess are caused by bacteria causing pulpal necrosis like Streptococci and Peptostreptococci which then progresses to formation of the dentoalveolar abscess. They can spread to the adjoining buccal, mandibular, submandibular, sublingual and submental spaces causing space infection and subsequently cellulitis formation<sup>13</sup>. Azodo et al assessed the presence of dentoalveolar abscess among Nigerian children and found that there was significant incidence of dentoalveolar abscess among children with deciduous first molar being the most commonly affected and untreated dental caries, the most common cause<sup>14</sup>.



### *Ankyloglossia*

Ankyloglossia is a congenital anomaly with an abnormally short lingual frenulum. It is also called as tongue tie with an 4.4% to 4.8% incidence in newborns and a male female ratio of 3:1.0. It can lead to difficulty in swallowing, sucking and speech in children<sup>15,16,17</sup>. The management includes frenectomy, lingual plasty and myofunctional training<sup>18</sup>.

### *White lesions*

The prevalence of white lesions in children include frictional keratosis, leukoedema and linea alba. They can present as ulcers, color changes, and alterations in size and configuration of oral anatomy. Discontinuation of causative habits and removal of the

causative irritant usually resolves the lesions<sup>19</sup>.

### *Eruption cyst*

The prevalence of eruption cyst is predominant in the Caucasian race. It is a soft tissue benign cyst associated with an erupting primary or permanent teeth and appears before the appearance of these teeth in the oral cavity<sup>20</sup>. They may disappear but should be treated if there is bleeding, pain or are infection. The management is drainage of the cystic contents<sup>21</sup>.

## **CONCLUSION**

A thorough oral examination plays an important role in the identification and





successful treatment of these oral lesions. Since oral and systemic health are interlinked, more emphasis has to be given on diagnosis and treatment of the same. Thus, this study plays a major important role in diagnosis of oral lesions which might go unnoticed in the initial stage. Therefore, a thorough and complete oral examination is mandatory in paediatric population to identify mucosal lesion and variation at an earlier level facilitating the management.

## REFERENCES

1. Unur M, Bektas Kayhan K, Altop MS, Boy Metin Z, Keskin Y. The prevalence of oral mucosal lesions in children:a single center study. J Istanbul Univ Fac Dent. 2015;49(3):29-38.
2. Yáñez, M.; Escobar,E.; Oviedo,C.; Stillfried, A.;Pennacchiotti, G. Prevalence of oral mucosal lesions in children. Int. J. Odontostomat., 10(3):463-468, 2016.
3. El Toum S, Cassia A, Bouchi N, Kassab I. Prevalence and Distribution of Oral Mucosal Lesions by Sex and Age Categories: A Retrospective Study of Patients Attending Lebanese School of Dentistry. Int J Dent.2018:4030134.
4. Wanda C. Gonsalves, Angela C. Chi, Brad W. Neville. Common Oral Lesions: Part I. Superficial Mucosal Lesions. Am Fam Physician. 2007 Feb 15;75(4):501-506.
5. Rioboo-Crespo MR, Planells-del Pozo P, Rioboo-García R. Epidemiology of the most common oral mucosal diseases in children.Med Oral Patol Oral Cir Bucal 2005;10:376-87.
6. Shulman JD. Prevalence of oral mucosal lesions in children and youths in the USA. Int J Paediatr Dent. 2005 Mar;15(2):89-97.



7. Hussein AA, Darwazeh AM, Al-Jundi SH. Prevalence of oral lesions among Jordanian children. Saudi J Oral Sci 2017;4:12-7
8. Ambika, L. & Keluskar, V. & Hugar, Shivayogim & Patil, S. Prevalence of oral mucosal lesions and variations in Indian public school children. Brazilian Journal of Oral Sciences. 2011;10. 288-293.
9. Mortazavi H, Safi Y, Baharvand M, Rahmani S. Diagnostic Features of Common Oral Ulcerative Lesions: An Updated Decision Tree. Int J Dent.2016;7278925.
10. Nandini DB, Bhavana SB, Deepak BS, Ashwini R. Paediatric Geographic Tongue: A Case Report, Review and Recent Updates. J Clin Diagn Res. 2016;10(2):ZE05-ZE9.
11. Pari A, Ilango P, Subbareddy V, Katamreddy V, Parthasarthy H. Gingival diseases in childhood - a review. J Clin Diagn Res. 2014;8(10):ZE01-ZE4.
12. Abrams RG, Romberg E. Gingivitis in children with malnutrition. The Journal of Clinical Pediatric Dentistry. 1999 ;23(3):189-194.
13. Farias HS, Coelho CSM, Costa CPA, Andrade CJ, Gonçalves FJ. Chronic dentoalveolar abscess in a pediatric patient with rare drainage. Rev Cubana Estomatol. 2019;56(4):1-13
14. Azodo CC, Chukwumah NM, Ezeja EB. Dentoalveolar abscess among children attending a dental clinic in Nigeria. Odontostomatol Trop. 2012 Sep;35(139):41-6.
15. Junqueira Marina Azevedo, Cunha Nayara Nery Oliveira, Costa e Silva Lidiane Lucas, Araújo Leandro Borges, Moretti Ana Beatriz Silveira, Couto Filho Carlos Eduardo Gomes et al . Surgical techniques for the treatment of ankyloglossia in children: a case series. J. Appl. Oral Sci. 2014 June ;22(3 ): 241-248.
16. Messner AH, Lalakea ML, Aby J, Macmahon J, Bair E. Ankyloglossia: incidence and associated feeding



difficulties. Arch Otolaryngol Head Neck Surg. 2000;126(1):36-9.

17. Friend GW, Harris EF, Mincer HH, Fong TL, Carruth KR. Oral anomalies in the neonate, by race and gender, in an urban setting. Pediatr Dent. 1990;12(3):157-61.

18. Ferrés-Amat E, Pastor-Vera T, Ferrés-Amat E, Mareque-Bueno J, Prats-Armengol J, Ferrés-Padró E. Multidisciplinary management of ankyloglossia in childhood. Treatment of 101 cases. A protocol. Med Oral Patol Oral Cir Bucal. 2016 Jan 1;21(1):e39-47.

19. Pinto, Andres & Haberland, Christel & Baker, Suher. (2014). Pediatric Soft Tissue Oral Lesions. Dental clinics of North America. 58. 437-453. 10.1016.

20. Nagaveni N B, Umashankara K V, Radhika N B, Maj Satisha T S. Eruption cyst: A literature review and four case reports. Indian J Dent Res 2011;22:148-51.

21. Dhawan P, Kochhar GK, Chachra S, Advani S. Eruption cysts: A series

of two cases. Dent Res J (Isfahan). 2012;9(5):647-650.