

Evaluation of the efficacy of Protaper, HERO SHAPER GOLD, ProTaper Universal retreatment, and R-Endo, for removing gutta-percha and AH plus sealer from root canals.

(Evaluación de la eficacia de Protaper, HERO SHAPER GOLD, retratamiento ProTaper Universal y R-Endo, para eliminar gutapercha y sellador AH plus de los conductos radiculares.)

Upendranatha Reddy N ¹, Gopi Krishna Reddy M ¹, Sujayeendranatha Reddy ¹E, Anusha Y ¹,
Nagalakshmi Reddy S ¹, Datta Sai Kiran K ¹

¹ G Pulla Reddy Dental College And Hospital, Kurnool, India

Received: 04 November 2024.

Accepted: 20 March 2025.

Online publication: 6 May 2025

[Original paper]

PPI: S2477-9369(24)13000X-OP

Abstract(english)

The purpose of the present study is to evaluate the efficacy of four different rotary NiTi files ProTaper files, HERO SHAPER GOLD files, ProTaper Universal retreatment files, and R-Endo files to remove GP and sealer from root canals with or without the use of passive ultrasonic irrigation using Irrisafe file under DOMS. This study hypothesizes that using PUI could result in better cleanliness of root canals after instrumentation for removal of GP and sealer. The study samples comprised 100 extracted single-rooted human maxillary anterior teeth and were collected from the Department of Oral and Maxillofacial Surgery, G. Pulla Reddy Dental College & Hospital, Kurnool. The t Test shows that there was a statistically significant difference between individual Sub groups of Groups I, II & IV ($p < 0.05$). No statistically significant difference between Subgroups of Group III ($p > 0.05$) but with percentage of remaining GP and sealer in the root canals after retreatment was comparatively greater in Subgroup A than in Sub group B. Under the experimental conditions, all the retreatment files left some amount of GP and sealer in the root canals and there was no significant difference between them. However, R-Endo is better following ProTaper Universal retreatment system proved, Protaper files, and HERO SHAPER GOLD files. Further use of passive ultrasonic irrigation with an Irrisafe file resulted in better cleanliness of the root canal wall after retreatment.

Keywords(english)

Endodontics, dental, files, irrigation, root canal, microscope.

Resumen(español)

El propósito del presente estudio es evaluar la eficacia de cuatro limas rotatorias NiTi diferentes: limas ProTaper, limas HERO SHAPER GOLD, limas de retratamiento ProTaper Universal y limas R-Endo para remover GP y sellador de conductos radiculares con o sin el uso de irrigación ultrasónica pasiva utilizando la lima Irrisafe bajo DOMS. Se hipotetizo de que el uso de PUI podría resultar en una mejor limpieza de los conductos radiculares después de la instrumentación para la remoción

de GP y sellador. Las muestras del estudio comprendieron 100 dientes anteriores maxilares humanos unirradiculares extraídos y se recolectaron del Departamento de Cirugía Oral y Maxilofacial, G. Pulla Reddy Dental College & Hospital, Kurnool. La prueba t muestra que hubo una diferencia estadísticamente significativa entre los subgrupos individuales de los Grupos I, II y IV ($p < 0,05$). No se observaron diferencias estadísticamente significativas entre los subgrupos del Grupo III ($p > 0,05$), pero el porcentaje de GP y sellador remanente en los conductos radiculares tras el retratamiento fue comparativamente mayor en el Subgrupo A que en el Subgrupo B. En conclusión, todas las limas de retratamiento dejaron cierta cantidad de GP y sellador en los conductos radiculares, sin que se observaran diferencias significativas entre ellas. Sin embargo, R-Endo es mejor tras el uso del sistema de retratamiento ProTaper Universal, las limas Protaper y las limas HERO SHAPER GOLD. El uso posterior de irrigación ultrasónica pasiva con una lima Irrisafe resultó en una mejor limpieza de la pared del conducto radicular tras el retratamiento.

Palabras clave(español)

Endodoncia, dental, limas, irrigación, conducto radicular, microscopio.

Introduction

Recently, rotary NiTi files specifically designed for removal of GP and other RC filling materials (1-7) have been introduced into the market, claiming rapid and effective in removal of RC filling material (8-13). Few of them are ProTaper Universal retreatment files (Dentsply Maillefer), R-Endo retreatment system (Micro Mega), Mtwo retreatment files (VDW, Munich, Germany), D-RaCe retreatment system (FKG Dentaire, La Chaux-de-Fonds, Switzerland). The use of passive ultrasonic irrigation (PUI) after instrumentation of RC has improved effect in removal of residual debris and smear layer (14). Purpose of the present study is to evaluate the efficacy of four different rotary NiTi files ProTaper files, HERO SHAPER GOLD files, ProTaper Universal retreatment files and R- Endo files to remove GP and sealer from root canals with or without use of passive ultrasonic irrigation using Irrisafe file under DOMS. Hypothesis of this study is that the use of PUI could result in better cleanliness of root canals after instrumentation for removal of GP and sealer.

Materiales y métodos

The present in vitro study was conducted in the department of Conservative dentistry and Endodontics, G. Pulla Reddy Dental College & Hospital, Kurnool, Andhra Pradesh. The study samples comprised of 100 extracted single rooted human maxillary anterior teeth and were collected from Department of Oral and Maxillofacial Surgery, G. Pulla Reddy Dental College & Hospital, Kurnool. The criteria for the selection of teeth were: Inclusion criteria: 1) Free of restorations. 2) With straight roots. 3) Complete root formation. Exclusion criteria: 1) Carious tooth. 2) Crack. 3) Fractured tooth.

Instruments: Micro motor hand piece. (NSK, Japan) Airotor hand piece. (NSK, Japan)Endo Access bur no. 1 (Dentsply Maillefer)Diamond discs. (Horico)Size 10, 15, 20 K file. (Mani)Endodontic torque controlled Rotary. (16:1 , X- Smart, Dentsply Maillefer)Warm thermo plasticizing obturation device. (E & Q plus, Meta Biomed, Korea)Hand pluggers. (Dentsply Maillefer)X ray machine .(Blux, Intra OS 70, Confident)Dental Operating microscope. (Labomed)Protaper rotary files. (Dentsply Maillefer)K3 XF files. (Sybron Endo)Protaper universal retreatment files. (Dentsply Maillefer)R-Endo files. (Micro Mega)Piezoelectronic unit. (Satelec, P5 Newtron XS)Ultrasonic endodontic file – Irrisafe File. (Satelec)Stereomicroscope. (Lynx, Lawrence & Mayo) Digital camera. (Nikon) [\[see supplementary material figure S1 and S2\]](#)

Materials used: 3% sodium hypochlorite. (Vishal Dento Care Pvt. Ltd.)Normal saline. (nirlife, Nirma limited)Distilled water. (Sreemanenterprise)Irrigation syringes and needles. (Ultradent)Paper points. (Meta Biomed) AH plus sealer. (Dentsply De Trey)Gutta-percha cones and pellets. (Dentsply Maillefer)Radiographs. (Carestream, E-Speed)Radiographic Developer & Fixer solution.CavitG. (3M ESPE)5% nitric acid. (SDFCL, SD Fine Chem Ltd.)Ethyl alcohol of 80%, 90%, 100%. (CS, Chinachangshu Yangyuan Chemicals)Methyl salicylate. (Himedia Laboratories Pvt Ltd.)

Specimen preparation: Teeth were stored in 3% sodium hypochlorite (NaOCL) for 24 hours to remove soft tissue debris and mechanically removed the calculus from tooth surface using ultrasonic scaler. Teeth were stored in distilled water until use.Access preparation was made on each tooth using high speed diamond bur using airotor hand piece with water coolant. A size 10 K-file was introduced into the canal until it was visible at the apical foramen. The working length was determined by subtracting 1mm from this

measurement. The crowns were decoronated with a diamond disk and straight handpiece to a standardized length to 16mm.

Root Canal Treatment: After establishment of glide path with no. 10 to no. 20 K files, RC biomechanical preparation was carried out with ProTaper rotary files as per manufacture recommendations. Root canals were shaped using S1 to reach working length followed by Sx for coronal flaring then S2 to reach working length. Followed by finishing of root canals using F1, F2, F3 to reach working length. At each change of instruments, root canals were irrigated using 28 gauge needles with 2 mL of 3% NaOCl. After completion of instrumentation, root canals were finally irrigated with 5 mL of normal saline and 5 mL of distilled water. The root canals were dried with paper points. Plugger was selected for each specimen which was taken to depth of approximately 3mm short of working length. Master cone was selected 1 to 2mm short of working length and was checked for tight apical tug back. Paste A and Paste B of AH plus sealer was mixed in equal proportion on mixing pad and coated to RC walls using paper points. Obturation was done with GP and AH plus sealer using continuous wave of condensation technique using E & Q plus. After placement of master cone in to RC, down pack was carried out using selected plugger attached to Pen of E & Q plus unit with continuous heat until plugger touches canal walls in the apical portion. Plugger was held in position for about 15 seconds to cool down the GP then again heat was activated to plugger for 1 second and withdrawn from the RC. Remaining portion of RC was back filled with thermo plasticized GP using Gun of E & Q plus unit set at 2000C. The injecting needle was positioned in the canal, preheated GP is then passively injected, the needle backs out of the canal and hand pluggers were used to compact the GP. The coronal access cavities were then sealed with Cavit G. Root canals of 100 teeth specimens after prepared and obturated they were radiographically evaluated in both mesio-distal (MD) and bucco-lingual (BL) direction for apical extent of obturation and any internal voids. Out of which 80 teeth presents better adaptation RC filling material with no internal voids and were selected for further retreatment. All 80 selected obturated teeth were stored at 100% humidity and 37°C for 30 days to allow complete sealer setting.

Retreatment Technique: The teeth were randomly divided into 4 groups with 20 specimens each. (n=20) Each group was divided in to two subgroups, A and B with 10 specimens each. Entire retreatment

procedure was performed under Dental Operating Microscope.

Group I – ProTaper Rotary File (DentsplyMaillefer). **Sub Group I A:** Rotary ProTaper NiTi files in an electric motor (X Smart), with a constant speed of 300 rpm were used with light apical pressure by the following sequence; Finishing files #3 (size 30, taper 0.09), #2 (size 25, taper 0.08), and #1 (size 20, taper 0.07) in a crown-down technique to remove the GP and sealer until the working length was reached. Finishing files #2 and #3 were used again to the working length to complete GP and sealer removal from the canal walls. **Sub Group I B:** Specimens of Sub Group I B were subjected to retreatment procedure with rotary ProTaper NiTi files as mentioned in Sub group I A along with passive ultrasonic irrigation was done with Irrisafe file (size 20) for 2 minutes at power setting “yellow 4” by Satelec, P5 Newtron XS piezoelectronic unit.

Group II – HERO SHAPER GOLD Rotary files. **Sub Group II A:** Rotary HERO SHAPER GOLD NiTi files with the electric motor (X Smart) at a constant speed of 300 rpm were used with a light apical pressure using the following sequence: Size 25 (taper 0.10), size 25 (0.08 taper), and size 20 (0.06 taper) in a crown-down technique to remove the GP and sealer until the working length was reached. Completion of GP removal and cleaning of canal walls was done using size 25 (0.06 taper) followed by size 30 (0.06 taper) to the working length. **Sub Group II B:** Specimens of Sub Group II B were subjected to retreatment procedure with rotary HERO SHAPER GOLD NiTi files as mentioned in Sub group II A along with passive ultrasonic irrigation was done with Irrisafe file (size 20) for 2 minutes at power setting “yellow 4” by Satelec, P5 Newtron XS piezoelectronic unit.

Group III- ProTaper Universal Rotary Retreatment files (DentsplyMaillefer). **Sub Group III A:** Rotary ProTaper Universal Retreatment files were used with an electric motor (X Smart) at a constant speed of 500 rpm. D1 with tip 30 and taper 0.09, D2 with tip 25 and taper 0.08 and D3 with tip 20 and taper 0.07 were used sequentially, applying a crown-down technique to remove GP and sealer, until the working length was reached. **Sub Group III B:** Specimens of Sub Group III B were subjected to retreatment procedure with rotary ProTaper Universal Retreatment NiTi files as mentioned in Sub group III A along with passive ultrasonic irrigation was done with Irrisafe file (size 20) for 2 minutes at power setting “yellow 4” by Satelec, P5 Newtron XS piezoelectronic unit.

Group IV - R-Endo System (Micro –Mega). Sub

Group IV A: Rotary R- Endo NiTi files were used for removal of GP and sealer with electric motor (X Smart) at a speed of 300 rpm. Rm stainless steel manual file (no.25, 4% taper) was used first to disrupt GP and center the NiTi files. It was followed by NiTi rotary files Re(no. 25, 12% taper), R1(no. 25, 8% taper), R2(no. 25, 6% taper), R3(no. 25, 4% taper) in crown down technique to reach working length. **Sub Group IV B:** Specimens of Sub Group IV B were subjected to retreatment procedure with rotary R-Endo NiTi files as mentioned in Sub group IV A along with passive ultrasonic irrigation was done with Irrisafe file (size 20) for 2 minutes at power setting “yellow 4” by Satelec, P5 Newtron XS piezoelectronic unit. The files were cleaned regularly using gauze to remove any obturated material and debris before being reintroduced in the root canal. Each file was discarded after being used in 5 specimens. During retreatment procedure Irrigation was performed with 28 28-gauge needle using 2 mL of 3% NaOCl at each instrument change. Retreatment was considered complete for all the groups when no filling material was observed on the instruments. Root canals were finally irrigated with 5 mL of normal saline and 5 mL of distilled water [\(see supplementary material Figure S3\)](#).

Evaluation of remaining gutta-percha and sealer: All the specimens were rendered transparent according to the clearing technique described by Don Robertson et al. Specimens were decalcified in 5 % nitric acid for 72 hours and then washed for 4 hours in running water and dehydrated in increasing concentrations of ethyl alcohol 80 % for 12 hours, 90 % for 1 hour, and 100% for 1 hour. The specimens were then cleared by placing in methyl salicylate solution until they become transparent. The amount of GP and sealer on the canal walls were estimated using

stereomicroscope by capturing images of transparent specimens in both MD and BL directions using digital camera at 8X magnification. The images were analyzed using Auto CAD 2004 software for the area of residual filling materials in square millimeters (mm²). Statistical

Analysis: All the data was analyzed using SPSS 21.0 version. Cleanliness of Root canals were analyzed using One way ANOVA with Turkeys multiple post-hoc test for Inter-group comparison and t test for Intra-group comparison. A p-value of < 0.05 was considered statistically significant.

Results

The obtained data were statistically analysed using One way ANOVA with Turkeys multiple post hoc test for inter Group comparison and t test for intra group comparison (P<0.05). In the present study four different rotary NiTi files ProTaper files HERO SHAPER GOLD, files, ProTaper Universal retreatment files and R-Endo files were used for removal of GP and sealer with or without use of PUI using Irrisafe file. The results for Canal Wall Cleanliness in present study have been discussed as follows

Inter group comparison between. Group I Vs. Group II / Group I Vs. Group III / Group I Vs. Group IV / Group II Vs. Group III / Group II Vs. Group IV / Group III Vs Group IV

Intra group comparison between. Sub Group I A Vs. Sub Group I B / Sub Group II A Vs. Sub Group II B / Sub Group III A Vs. Sub Group III B / Sub Group IV A Vs. Sub Group IV B

From table 1 and table 2 following data were analysed. i. Group I has a mean percentage of remaining guttapercha and sealer of about 28.84% when specimens viewed in MD direction and of about

Table. 1. Inter Group Comparison of Mesio Distal specimens by one way ANOVA.

Group	Coronal		Middle		Apical		Total	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD
Group I	25.81	16.67	27.90	26.62	28.62	17.40	28.84	15.35
Group II	26.09	18.83	30.84	24.58	26.67	14.50	27.86	15.14
Group III	18.04	15.46	28.78	26.35	32.56	16.24	23.14	13.55
Group IV	14.94	15.96	24.92	23.59	31.51	20.22	21.81	10.65
F- Value	2.2348		0.1891		0.4882		1.2574	
p-value	0.0910		0.9035		0.6915		0.2951	
Pair wise comparison of groups by Tukeys multiple posthoc procedures								
Group I Vs II	P=0.9999		p=0.9830		p=0.9843		p=0.9961	
Group I Vs III	p=0.4641		p=0.9996		p=0.8875		p=0.5625	
Group I Vs IV	p=0.1795		p=0.9822		p=0.9512		p=0.3782	
Group II Vs III	p=0.4324		p=0.9940		p=0.7023		p=0.7018	
Group II Vs IV	p=0.1618		p=0.8804		p=0.8106		p=0.5106	
Group III Vs IV	p=0.9364		p=0.9626		p=0.9975		p=0.9900	

*p < 0.05

Table. 2. Inter Group Comparison of Bucco Lingual specimens by one way ANOVA.

Group	Coronal		Middle		Apical		Total	
	Mea n	SD	Mean	SD	Mean	SD	Mean	SD
Group I	21.74	16.10	22.64	25.98	25.74	18.48	22.94	16.71
Group II	24.50	18.02	30.10	25.45	25.07	17.54	26.31	16.02
Group III	16.67	15.50	25.23	23.83	22.30	16.98	20.95	13.87
Group IV	9.04	10.34	24.45	25.57	29.66	20.19	17.49	10.93
F- Value	3.9611		0.3191		0.5476		1.2876	
p-value	0.0111*		0.8115		0.6513		0.2848	
Pair wise comparison of groups by Tukeys multiple posthoc procedures								
Group I Vs II	p=0.9397		p=0.7862		p=0.9995		p=0.8839	
Group I Vs III	p=0.7209		p=0.9882		p=0.9341		p=0.9728	
Group I Vs IV	p=0.0495*		p=0.9959		p=0.9059		p=0.6382	
Group II Vs III	p=0.3719		p=0.9284		p=0.9639		p=0.6509	
Group II Vs IV	p=0.0105*		p=0.8937		p=0.8583		p=0.2296	
Group III Vs IV	p=0.3950		p=0.9997		p=0.5855		p=0.8754	

*p < 0.05

22.94% when specimens viewed in BL direction. **ii.** Group II has a mean percentage of remaining guttapercha and sealer of about 27.86% when specimens viewed in MD direction and of about 26.31% when specimens viewed in BL direction. **iii.** Group II has

a mean percentage of remaining guttapercha and sealer of about 23.14% when specimens viewed in MD direction and of about 20.95% when specimens viewed in BL direction. **iv.** Group II has a mean percentage of remaining guttapercha and sealer of about 21.81% when specimens viewed in MD direction and of about 17.49% when specimens viewed in BL direction. **v.** The percentage of remaining guttapercha and sealer in coronal, middle and apical thirds' were comparative more in middle and apical thirds' than in coronal thirds'. No significance difference in coronal, middle and apical thirds compared to other groups. But Significance difference between Group I vs IV = $p=0.0495$, Group II vs IV = $p=0.0105$ (Specimens viewed in BL direction) observed in respect to coronal third (Table 2)

Inter group comparison. Table 1 & 2, Figure 1 & 2) Using One way ANOVA with Tukeys multiple post hoc test. **i.** Comparison between Group I and Group II: There was no statistical significance difference ($p>0.05$) between Group I and Group II specimens viewed in MD & BL




Figure 1. Inter Group Comparison of Mesio Distal specimens




Figure 2. Inter Group Comparison of Bucco Lingual specimens




Figure 3. Intra group comparison of Group I

Table. 3. Intra Group Comparison of Group I by t test.


		Sub group A		Sub group B		t-value	p-value
		Mean	SD	Mean	SD		
Mesio Distal	Coronal	31.61	14.10	20.01	17.70	1.6218	0.1222
	Middle	38.63	32.45	17.18	13.68	1.9264	0.0700
	Apical	37.24	17.82	20.00	12.51	2.5039	0.0221*
	Total	36.54	14.81	21.14	12.11	2.5448	0.0203*
Bucco Lingual	Coronal	23.76	15.15	19.71	17.58	0.5521	0.5877
	Middle	33.83	32.66	11.46	8.97	2.0991	0.0500*
	Apical	36.31	20.11	15.17	8.29	3.0727	0.0066*
	Total	29.21	20.12	16.68	9.86	1.7688	0.0491*

*p < 0.05

direction with $p=0.9961$ & 0.8839 respectively. **ii.** Comparison between Group I and Group III: There was no statistical significance difference ($p>0.05$) between Group I and Group III specimens viewed in MD & BL direction with $p=0.5625$ & 0.9728 respectively. **iii.** Comparison between Group I and Group IV: There was no statistical significance difference ($p>0.05$) between Group I and Group IV specimens viewed in MD & BL direction with $p=0.3782$ & 0.6382 respectively. **iv.** Comparison between Group II and Group III: There was no statistical significance difference ($p>0.05$) between Group II and Group III specimens viewed in MD & BL direction with $p=0.7018$ & 0.6509 respectively. **v.** Comparison between Group II and Group IV: There was no statistical significance difference ($p>0.05$) between Group II and Group IV specimens viewed in MD & BL direction with $p=0.5106$ & 0.2296 respectively. **vi.** Comparison between Group III and Group IV: There was no statistical significance difference ($p>0.05$) between Group III and Group IV specimens viewed in MD & BL direction with $p=0.9900$ & 0.8754 respectively.

Intra group comparison. Using t test. i. Sub Group I A and Sub Group I B: (Table 3, Figure 3) There was statistical significance difference ($p<0.05$) between Sub Group I A and Sub Group I B specimens viewed in MD & BL direction with $p=0.0203$ & 0.0491 respectively. **ii.** Sub Group II A and Sub Group II B: (Table 4, Figure 4) There was statistical significance difference ($p<0.05$) between Sub Group II A and Sub Group II B specimens viewed in

MD & BL direction with $p=0.4841$ & 0.0165 respectively. **iii.** Sub Group III A and Sub Group III B: (Table 5, Figure 5) There was statistical no significance difference ($p>0.05$) between Sub Group III A and Sub Group III B specimens viewed in MD & BL direction with $p=0.6242$ & 0.5541 respectively. **iv.** Sub Group IV A and Sub Group IV B: (Table 6, Figure 6) There was statistical significance difference ($p<0.05$) between Sub Group IV A and Sub Group IV B specimens viewed in MD & BL direction with $p=0.0074$ & 0.0201 respectively. None of the retreatment techniques completely removed the root canal filling material from root canals. One way ANOVA

**Figure 4. Intra group Comparison of Group II****Table. 4. Intra Group Comparison of Group II by t test.**

		Sub group A		Sub group B		t-value	p-value
		Mean	SD	Mean	SD		
Mesio Distal	Coronal	23.34	20.08	28.84	18.12	-0.6429	0.5284
	Middle	46.45	20.33	15.24	17.91	3.6425	0.0019*
	Apical	33.73	16.64	19.62	7.49	2.4469	0.0249*
	Total	32.08	15.70	23.64	14.06	1.2666	0.4841*
Bucco Lingual	Coronal	26.21	20.01	22.80	16.70	0.4136	0.6841
	Middle	48.95	20.57	11.25	12.42	4.9631	0.0001*
	Apical	35.10	18.12	15.04	9.89	3.0727	0.0066*
	Total	34.57	16.48	18.05	10.91	2.6435	0.0165*

*p < 0.05

Table. 5. Intra Group Comparison of Group III by t test.

		Sub group A		Sub group B		t-value	p-value
		Mean	SD	Mean	SD		
Mesio Distal	Coronal	13.83	11.81	22.25	18.04	-1.2359	0.2324
	Middle	37.91	22.30	19.66	27.99	1.6122	0.1243
	Apical	38.74	14.96	26.38	15.75	1.7997	0.0887
	Total	24.68	11.10	21.60	16.09	0.4985	0.6242
Bucco Lingual	Coronal	14.84	13.76	18.51	17.61	-0.5196	0.6097
	Middle	32.04	21.81	18.42	24.90	1.3012	0.2096
	Apical	29.25	17.69	15.35	13.73	1.9630	0.0653
	Total	22.85	12.20	19.05	15.78	0.6030	0.5541

*p < 0.05

with Turkey's multiple post hoc test shows that there was no statistical significance difference between the groups ($p > 0.05$), but with difference in the mean values. The order of Groups with lesser to higher mean percentage of remaining GP and sealer in the root canals after retreatment was as follows Group IV, Group III, Group I & Group II. ($\text{Group IV} < \text{III} < \text{I} < \text{II}$). The difference in the mean percentage of remaining GP and sealer in the root canals after retreatment between Group II and Group IV is about 6.05% when viewed in MD direction and 8.82% when viewed in BL direction. Test shows that there was statistical significance difference between individual Sub groups of Groups I, II & IV ($p < 0.05$). And no statistical significance difference between Subgroups of Group III ($p > 0.05$) but with percentage of remaining GP and sealer in the root canals after retreatment was comparatively greater in Sub group A than in Sub group B.

Discussion

The present in vitro study was done to compare the efficacy of rotary NiTi files Protaper files, files and retreatment systems ProTaper Universal retreatment system and R- Endo retreatment system in

removal of gutta-percha and AH plus sealer with or without use of passive ultrasonic irrigation with Irrisafe file, under Dental Operating Microscope and subjects were evaluated using clearing technique under the stereomicroscope and photographs (4,6,7,20,23,25,35,65,68,69,70,71,72). In the present study RC specimens were obturated using continuous wave of condensation. (E & Q plus) (3,4,6-10,15-23,30,31,34-36,39,41,43,44,48,50,51,54,57,73,74,75,76,77).

Group I Vs. II: MD viewed specimens ($p = 0.9961$) & BL viewed specimens ($p = 0.8839$); Group I Vs. III: MD viewed specimens ($p = 0.5625$) & BL viewed specimens ($p = 0.9728$); Group I Vs. IV: MD viewed specimens ($p = 0.3782$) & BL viewed specimens ($p = 0.6382$); Group II Vs. III: MD viewed specimens ($p = 0.7018$) & BL viewed specimens ($p = 0.6509$); Group II Vs. IV: MD viewed specimens ($p = 0.5106$) & BL viewed specimens ($p = 0.2296$); Group III Vs. IV: MD viewed specimens ($p = 0.9900$) & BL viewed specimens ($p = 0.8754$).

Though there was no significance difference between the groups, the order of sequence of groups with less to higher left over residual GP and sealer in RCs is Group IV- R – Endo retreatment system, Group III- Protaper universal retreatment system, Group I-




Figure 5. Intra group Comparison of Group III




Figure 6. Intra group Comparison of Group IV

Table. 6. Intra Group Comparison of Group IV by t test.

		Sub group A		Sub group B		t-value	p-value
		Mean	SD	Mean	SD		
Mesio Distal	Coronal	21.79	15.02	8.08	14.41	2.0899	0.0500*
	Middle	25.94	24.69	23.89	23.72	0.1901	0.8514
	Apical	42.01	16.80	21.02	18.34	2.6704	0.0156*
	Total	27.82	10.21	15.79	7.40	3.0157	0.0074*
Bucco Lingual	Coronal	13.22	11.83	4.86	6.86	1.9339	0.0690
	Middle	27.93	28.49	20.98	23.28	0.5979	0.5574
	Apical	41.66	18.28	17.66	14.39	3.2622	0.0043*
	Total	22.98	12.03	12.00	6.36	2.5510	0.0201*

*p < 0.05

Protaper files and Group II – HERO SHAPER GOLD files. There was a significance ($p < 0.05$) difference in between the individual group that is intra group comparison, with or without the use of passive ultrasonic irrigation. Sub Group I A Vs. Sub Group I B: MD viewed specimens ($p = 0.0203^*$) & BL viewed specimens ($p = 0.0491^*$); Sub Group II A Vs. Sub Group II B: MD viewed specimens ($p = 0.4841^*$) & BL viewed specimens ($p = 0.0165^*$); Sub Group III A Vs. Sub Group III B: MD viewed specimens ($p = 0.6242$) & BL viewed specimens ($p = 0.5541$); Sub Group IV A Vs. Sub Group IV B: MD viewed specimens ($p = 0.0074^*$) & BL viewed specimens ($p = 0.0201^*$). Remaining filling material distribution in coronal, middle and apical thirds of root canals, much of remaining filling material was observed in the middle and apical thirds. The reason could be because in most of present retreatment techniques followed crown down technique for removal of GP and sealer and files used for coronal third of root canals have greater taper than used for middle and apical thirds. But there is no significance difference in distribution of remaining filling materials in respect to thirds in all the groups with $p > 0.05$. From results of present study it was shown that the residual GP and sealer are more in the specimens when viewed in BL direction. This is due to the fact that though RCs were standardized with same biomechanical preparation in all specimens, RCs of maxillary anteriors are more or less oval in shape and files used for retreatment purpose are mostly round in shape. R-Endo files are comparatively effective than ProTaper Universal retreatment files, ProTaper files and HERO SHAPER GOLD Files. The reason could be in R – Endo system it was provided Re file with a taper of 0.12, tip size 25 of 10 mm length. It has aggressive cutting edges and aid in removal of root canal filling material. Hence in the R – Endo group presents lesser filling material in coronal and middle thirds compared to other groups. Significance difference between Group I vs IV = $p = 0.0495$, Group II vs IV = $p = 0.0105$ observed in respect to coronal third. But there is no significance difference

in middle and apical thirds compared to other groups. The results showed that no significant difference was observed between the filling materials on terms of their removal. Manual instrumentation left more filling debris on the root canal walls when compared to HERO SHAPER GOLD and ProTaper.8 Results showed that there was no statistically significant difference among the others techniques: ProFile, ProTaper and HERO SHAPER GOLD when compared with GT. Rotary files GT, ProFile, ProTaper and HERO SHAPER GOLD were more effective in removing gutta-percha than manual and Hero instruments (44). The results showed that no significant differences were observed between the rotary systems in terms of the area of filling material left within the canals. There were statistically significant differences between the filling materials: Mtwo Retreatment files were more rapid when removing filling material than ProTaper Retreatment files and Twisted Files (43). Results showed that all instrumentation techniques left gutta-percha and sealer remnants inside the root canals. R-Endo instrumentation was significantly more effective ($P < 0.05$) than MTwo retreatment files in removing gutta-percha from the middle and apical thirds (48). Results of this study can be correlate with present study where R – Endo is showed as efficient compared to others retreatment techniques. ProTaper Universal rotary retreatment system without chloroform was faster and effective (37). Most remnants were found in the apical third of the canals.46Which is comparable to present study that most of filling material left was mostly present in apical third of the canals. In present study AH plus is used as root canal sealer and present more root canal filling debris than other studies where zinc oxide and calcium hydroxide based sealers are used (21, 23,46). Results showed that remaining filling material was observed in all specimens. The mean volume of remaining material was higher in the continuous wave of condensation groups than in the cold lateral condensation groups, especially in the apical portions of the root canals (58,72). The results showed significant

differences between the two removal techniques. Gutta-percha was more efficiently removed by using hand K-files compared to ProTaper universal retreatment files. Reason explained for this finding was that all canals were enlarged to a size F3 ProTaper file, which has a tip size of 30 and 9% taper, whereas the D3 ProTaper retreatment file has a tip size of 20 and 7% taper, which meant the D3 file tip did not engage with the canal walls (6). However, the high degree of filling material remaining in this study could be because of the constant size of retreatment files (size 25) rather than the instruments used during root canal preparation (size 30). Further enlargement of root canals beyond the canal dimension at the time of removal of root filling could have resulted in a significant reduction in material and in cleaner walls (39). This study was correlative to present study in the aspect of method of evaluation of residual RC filling material using clearing technique. And after removal of RC filling material, further RCs were instrumented with Protaper rotary instruments. Results showed ProTaper Universal rotary retreatment system and with further canal reparation accomplished with ProTaper rotary comparatively left less residual GP (34). The results showed that residues present after the use of the ProTaper Universal rotary files is comparatively more than following the supplementary application of the SAF. It was concluded that the use of the SAF after rotary instrumentation using ProTaper Universal retreatment files resulted in a significant reduction in the amount of filling residue in curved canals of mandibular molars (50). Results revealed that ultrasound/xylol led to lower percentages of remaining sealer, significantly different from the Protaper retreatment, Protaper retreatment /xylol and ultrasound which were similar. Ultrasound/xylol led to significantly lower percentages of remaining sealer on

the canal walls when compared to other groups.65 these results can be correlate with present study where PUI was used in presence of NaoCl instead of RC solvents, Xylol.64 Results showed that there were no significant differences between the groups or among the root canal thirds within each group. PUI with Endosolv R was not effective in the removal of filling debris from root canal walls (14). These results supports the present study in which passive ultrasonic instrumentation has negative out come during root canal retreatment with the use of RC solvents, hence instead of RC solvents NaoCl was used during PUI.. The results showed average percentage of remaining gutta-percha/sealer was higher in convetional technique than convetional technique in combination with burs, solvent, ultrasonics plus clinical operating microscope showing a statistically significant difference. The use of the DOMS and ultrasonic tips removed the filling material from root canal walls better (12). The root canal cleanliness is best achieved when retreatment is performed under a DOMS (53). The results of present study demonstrate that under the experimental conditions, all the retreatment files left some amount of GP and sealer in the root canals and there was no significant difference between them.

In conclusion, the R – Endo retreatment system and ProTaper Universal rotary retreatment system have advantages over other retreatment files. No need of solvents, minimizes smearing of GP and sealer on RC walls. Time saving or faster. Instrument design specially designed for retreatment of root canals.

Conflicts of interest

None to declare

Referencias

1. Cohen S, Hargreaves KM. Pathways of the pulp. 10th ed. Atlanta: Mosby:2005. Roda RS, Gettleman BH. Chapter 25. Nonsurgical endodontic retreatment. 944-1010.
2. Friedman S, Abitbol S, Lawrence HP. Treatment outcome in endodontics: the Toronto Study. Phase 1: initial treatment. J Endod. 2003 Dec;29: 787-93. [\[PubMed\]](#) [\[Google Scholar\]](#)
3. Torabinejad M, Corr R, Handysides R, Shabahang S. Outcomes of nonsurgical retreatment and endodontic surgery: a systematic review. J Endod. 2009; 35: 930-7. [\[PubMed\]](#) [\[Google Scholar\]](#)
4. Saad AY, Al-Hadlaq SM, Al-Katheeri NH. Efficacy of two rotary NiTi instruments in the removal of Gutta-Percha during root canal retreatment. J Endod. 2007;33: 38-41. [\[PubMed\]](#) [\[Google Scholar\]](#)
5. Pirani C, Pelliccioni GA, Marchionni S, Montebugnoli L, Piana G, Prati C. Effectiveness of three different retreatment techniques in canals filled with compacted gutta-percha or Thermafil: a scanning electron microscope study. J Endod. 2009; 35:1433-40. [\[PubMed\]](#) [\[Google Scholar\]](#)
6. Hammad M, Qualtrough A, Silikas N. Three-dimensional evaluation of effectiveness of hand and rotary instrumentation for retreatment of canals filled with different materials. J Endod. 2008; 34: 1370-3. [\[PubMed\]](#) [\[Google Scholar\]](#)
7. Patil SA, Dodwad PK, Patil AA. An in vitro comparison of bond strengths of Gutta-percha/AH Plus, Resilon/Epiphany self-

- etch and EndoREZ obturation system to intraradicular dentin using a push-out test design. J Conserv Dent. 2013; 16: 238-42. [\[PubMed\]](#) [\[Google Scholar\]](#)
8. de Carvalho Maciel AC, Zaccaro Scelza MF. Efficacy of automated versus hand instrumentation during root canal retreatment: an ex vivo study. Int Endod J. 2006; 39: 779-84. [\[PubMed\]](#) [\[Google Scholar\]](#)
9. Horvath SD, Altenburger MJ, Naumann M, Wolkewitz M, Schirrmeister JF. Cleanliness of dentinal tubules following gutta-percha removal with and without solvents: a scanning electron microscopic study. Int Endod J. 2009; 42: 1032-8. [\[PubMed\]](#) [\[Google Scholar\]](#)
10. Mushtaq M, Farooq R, Ibrahim M, Khan FY. Dissolving efficacy of different organic solvents on gutta-percha and resilon root canal obturating materials at different immersion time intervals. J Conserv Dent. 2012; 15: 141-5. [\[PubMed\]](#) [\[Google Scholar\]](#)
11. Só MV, Saran C, Magro ML, Vier-Pelisser FV, Munhoz M. Efficacy of ProTaper retreatment system in root canals filled with gutta-percha and two endodontic sealers. J Endod. 2008; 34: 1223-5. [\[PubMed\]](#) [\[Google Scholar\]](#)
12. de Mello Junior JE, Cunha RS, Bueno CE, Zuolo ML. Retreatment efficacy of gutta-percha removal using a clinical microscope and ultrasonic instruments: part I--an ex vivo study. Oral Surg Oral Med Oral Pathol Oral Radiol Endod. 2009;108: e59-62. [\[PubMed\]](#) [\[Google Scholar\]](#)
13. Unal GC, Kaya BU, Taç AG, Keçeci AD. A comparison of the efficacy of conventional and new retreatment instruments to remove gutta-percha in curved root canals: an ex vivo study. Int Endod J. 2009; 42: 344-50. [\[PubMed\]](#) [\[Google Scholar\]](#)
14. Müller GG, Schönhofen ÂP, Móra PM, Grecca FS, Só MV, Bodanezi A. Efficacy of an organic solvent and ultrasound for filling material removal. Braz Dent J. 2013; 24: 585-90. [\[PubMed\]](#) [\[Google Scholar\]](#)
15. Wilcox LR. Endodontic retreatment: ultrasonics and chloroform as the final step in reinstrumentation. J Endod. 1989; 15: 125-8. [\[PubMed\]](#) [\[Google Scholar\]](#)
16. Ladley RW, Campbell AD, Hicks ML, Li SH. Effectiveness of halothane used with ultrasonic or hand instrumentation to remove gutta-percha from the root canal. J Endod. 1991; 17:221-4. [\[PubMed\]](#) [\[Google Scholar\]](#)
17. Hülsmann M, Stotz S. Efficacy, cleaning ability and safety of different devices for gutta-percha removal in root canal retreatment. Int Endod J. 1997; 30: 227-33. [\[PubMed\]](#) [\[Google Scholar\]](#)
18. Frajlich SR, Goldberg F, Massone EJ, Cantarini C, Artaza LP. Comparative study of retreatment of Thermafil and lateral condensation endodontic fillings. Int Endod J. 1998; 31: 354-7. [\[PubMed\]](#) [\[Google Scholar\]](#)
19. Imura N, Kato AS, Hata GI, Uemura M, Toda T, Weine F. A comparison of the relative efficacies of four hand and rotary instrumentation techniques during endodontic retreatment. Int Endod J. 2000; 33: 361-6. [\[PubMed\]](#) [\[Google Scholar\]](#)
20. Ferreira JJ, Rhodes JS, Ford TR. The efficacy of gutta-percha removal using Profiles. Int Endod J. 2001; 34: 267-74. [\[PubMed\]](#) [\[Google Scholar\]](#)
21. Betti LV, Bramante CM. Quantec SC rotary instruments versus hand files for gutta-percha removal in root canal retreatment. Int Endod J. 2001; 34: 514-9. [\[PubMed\]](#) [\[Google Scholar\]](#)
22. Hülsmann M, Bluhm V. Efficacy, cleaning ability and safety of different rotary NiTi instruments in root canal retreatment. Int Endod J. 2004; 37: 468-76. [\[PubMed\]](#) [\[Google Scholar\]](#)
23. Masiero AV, Barletta FB. Effectiveness of different techniques for removing gutta-percha during retreatment. Int Endod J. 2005; 38: 2-7. [\[PubMed\]](#) [\[Google Scholar\]](#)
24. Kosti E, Lambrianidis T, Economides N, Neofitou C. Ex vivo study of the efficacy of H-files and rotary Ni-Ti instruments to remove gutta-percha and four types of sealer. Int Endod J. 2006; 39: 48-54. [\[PubMed\]](#) [\[Google Scholar\]](#)
25. Schirrmeister JF, Meyer KM, Hermanns P, Altenburger MJ, Wrbas KT. Effectiveness of hand and rotary instrumentation for removing a new synthetic polymer-based root canal obturation material (Epiphany) during retreatment. Int Endod J. 2006; 39: 150-6. [\[PubMed\]](#) [\[Google Scholar\]](#)
26. de Oliveira DP, Barbizam JV, Trope M, Teixeira FB. Comparison between gutta-percha and resilon removal using two different techniques in endodontic retreatment. J Endod. 2006; 32: 362-4. [\[PubMed\]](#) [\[Google Scholar\]](#)
27. Zmner O, Pameijer CH, Banegas G. Retreatment efficacy of hand versus automated instrumentation in oval-shaped root canals: an ex vivo study. Int Endod J. 2006; 39: 521-6. [\[PubMed\]](#) [\[Google Scholar\]](#)
28. Schirrmeister JF, Hermanns P, Meyer KM, Goetz F, Hellwig E. Detectability of residual Epiphany and gutta-percha after root canal retreatment using a dental operating microscope and radiographs--an ex vivo study. Int Endod J. 2006; 39: 558-65. [\[PubMed\]](#) [\[Google Scholar\]](#)
29. Barletta FB, Rahde Nde M, Limongi O, Moura AA, Zanesco C, Mazocatto G. In vitro comparative analysis of 2 mechanical techniques for removing gutta-percha during retreatment. J Can Dent Assoc. 2007; 73: 65. [\[PubMed\]](#) [\[Google Scholar\]](#)
30. Taşdemir T, Er K, Yildirim T, Celik D. Efficacy of three rotary NiTi instruments in removing gutta-percha from root canals. Int Endod J. 2008 Mar;41: 191-6. [\[PubMed\]](#) [\[Google Scholar\]](#)
31. Zanettini PR, Barletta FB, de Mello Rahde N. In vitro comparison of different reciprocating systems used during endodontic retreatment. Aust Endod J. 2008; 34: 80-5. [\[PubMed\]](#) [\[Google Scholar\]](#)
32. Cunha RS, De Martin AS, Barros PP, da Silva FM, Jacinto RC, Bueno CE. In vitro evaluation of the cleansing working time and analysis of the amount of gutta-percha or Resilon remnants in the root canal walls after instrumentation for endodontic retreatment. J Endod. 2007; 33: 1426-8. [\[PubMed\]](#) [\[Google Scholar\]](#)
33. Barletta FB, de Sousa Reis M, Wagner M, Borges JC, Dall'Agnol C. Computed tomography assessment of three techniques for removal of filling material. Aust Endod J. 2008; 34: 101-5. [\[PubMed\]](#) [\[Google Scholar\]](#)
34. Gu LS, Ling JQ, Wei X, Huang XY. Efficacy of ProTaper Universal rotary retreatment system for gutta-percha removal from root canals. Int Endod J. 2008 ;41: 288-95. [\[PubMed\]](#) [\[Google Scholar\]](#)
35. Giuliani V, Cocchetti R, Pagavino G. Efficacy of ProTaper universal retreatment files in removing filling materials during root canal retreatment. J Endod. 2008; 34: 1381-4. [\[PubMed\]](#) [\[Google Scholar\]](#)

36. Aydin B, Köse T, Calışkan MK. Effectiveness of HERO 642 versus Hedström files for removing gutta-percha fillings in curved root canals: an ex vivo study. *Int Endod J.* 2009; 42: 1050-6. [\[PubMed\]](#) [\[Google Scholar\]](#)
37. Takahashi CM, Cunha RS, de Martin AS, Fontana CE, Silveira CF, da Silveira Bueno CE. In vitro evaluation of the effectiveness of ProTaper universal rotary retreatment system for gutta-percha removal with or without a solvent. *J Endod.* 2009; 35: 1580-3. [\[PubMed\]](#) [\[Google Scholar\]](#)
38. Uezu MK, Britto ML, Nabeshima CK, Pallotta RC. Comparison of debris extruded apically and working time used by ProTaper Universal rotary and ProTaper retreatment system during gutta-percha removal. *J Appl Oral Sci.* 2010; 18: 542-5. [\[PubMed\]](#) [\[Google Scholar\]](#)
39. Fenoul G, Meless GD, Pérez F. The efficacy of R-Endo rotary NiTi and stainless-steel hand instruments to remove gutta-percha and Resilon. *Int Endod J.* 2010; 43: 135-41. [\[PubMed\]](#) [\[Google Scholar\]](#)
40. Roggendorf MJ, Legner M, Ebert J, Fillery E, Frankenberger R, Friedman S. Micro-CT evaluation of residual material in canals filled with Activ GP or GuttaFlow following removal with NiTi instruments. *Int Endod J.* 2010; 43: 200-9. [\[PubMed\]](#) [\[Google Scholar\]](#)
41. Bramante CM, Fidelis NS, Assumpção TS, Bernardineli N, Garcia RB, Bramante AS, de Moraes IG. Heat release, time required, and cleaning ability of MTwo R and ProTaper universal retreatment systems in the removal of filling material. *J Endod.* 2010; 36: 1870-3. [\[PubMed\]](#) [\[Google Scholar\]](#)
42. Ramzi H, Shokouhinejad N, Saghiri MA, Samieefard A. Efficacy of Three Different Methods in the Retreatment of Root Canals Filled with Resilon/Epiphany SE. *Iran Endod J.* 2010; 5: 161-6. [\[PubMed\]](#) [\[Google Scholar\]](#)
43. Marfisi K, Mercade M, Plotino G, Duran-Sindreu F, Bueno R, Roig M. Efficacy of three different rotary files to remove gutta-percha and Resilon from root canals. *Int Endod J.* 2010; 43: 1022-8. [\[PubMed\]](#) [\[Google Scholar\]](#)
44. Fariniuk LF, Westphalen VP, Silva-Neto UX, Carneiro E, Baratto Filho F, Fidel SR, Fidel RA. Efficacy of five rotary systems versus manual instrumentation during endodontic retreatment. *Braz Dent J.* 2011; 22: 294-8. [\[PubMed\]](#) [\[Google Scholar\]](#)
45. Reddy S, Neelakantan P, Saghiri MA, Lotfi M, Subbarao CV, Garcia-Godoy F, Gutmann JL. Removal of gutta-percha/zinc-oxide-eugenol sealer or gutta-percha/epoxy resin sealer from severely curved canals: an in vitro study. *Int J Dent.* 2011; 2011: 541831. [\[PubMed\]](#) [\[Google Scholar\]](#)
46. Dadresanfar B, Mehrvarzfar P, Saghiri MA, Ghafari S, Khalilak Z, Vatanpour M. Efficacy of two rotary systems in removing gutta-percha and sealer from the root canal walls. *Iran Endod J.* 2011 Spring; 6: 69-73. [\[PubMed\]](#) [\[Google Scholar\]](#)
47. Siotia J, Acharya SR, Gupta SK. Efficacy of ProTaper Retreatment System in Root Canals Obturated with Gutta-Percha Using Two Different Sealers and GuttaFlow. *Int J Dent.* 2011;2011: 676128. [\[PubMed\]](#) [\[Google Scholar\]](#)
48. Mollo A, Botti G, Principi Goldoni N, Randellini E, Paragliola R, Chazine M, Ounsi HF, Grandini S. Efficacy of two Ni-Ti systems and hand files for removing gutta-percha from root canals. *Int Endod J.* 2012; 45: 1-6. [\[PubMed\]](#) [\[Google Scholar\]](#)
49. Kfir A, Tsisis I, Yakirevich E, Matalon S, Abramovitz I. The efficacy of five techniques for removing root filling material: microscopic versus radiographic evaluation. *Int Endod J.* 2012; 45: 35-41. [\[PubMed\]](#) [\[Google Scholar\]](#)
50. Abramovitz I, Relles-Bonar S, Baransi B, Kfir A. The effectiveness of a self-adjusting file to remove residual gutta-percha after retreatment with rotary files. *Int Endod J.* 2012; 45: 386-92. [\[PubMed\]](#) [\[Google Scholar\]](#)
51. Rödiger T, Hausdörfer T, Konietzschke F, Dullin C, Hahn W, Hülsmann M. Efficacy of D-RaCe and ProTaper Universal Retreatment NiTi instruments and hand files in removing gutta-percha from curved root canals - a micro-computed tomography study. *Int Endod J.* 2012; 45: 580-9. [\[PubMed\]](#) [\[Google Scholar\]](#)
52. Jayasenthil A, Sathish ES, Prakash P. Evaluation of manual and two-rotary niti retreatment systems in removing gutta-percha obturated with two root canal sealers. *ISRN Dent.* 2012;2012: 208241. [\[PubMed\]](#) [\[Google Scholar\]](#)
53. Chauhan R, Tikku A, Chandra A. Detection of residual obturation material after root canal retreatment with three different techniques using a dental operating microscope and a stereomicroscope: An in vitro comparative evaluation. *J Conserv Dent.* 2012; 15: 218-22. [\[PubMed\]](#) [\[Google Scholar\]](#)
54. Akhavan H, Azdadi YK, Azimi S, Dadresanfar B, Ahmadi A. Comparing the Efficacy of Mtwo and D-RaCe Retreatment Systems in Removing Residual Gutta-Percha and Sealer in the Root Canal. *Iran Endod J.* 2012 Summer; 7: 122-6. [\[PubMed\]](#) [\[Google Scholar\]](#)
55. Solomonov M, Paqué F, Kaya S, Adigüzel O, Kfir A, Yiğit-Özer S. Self-adjusting files in retreatment: a high-resolution micro-computed tomography study. *J Endod.* 2012; 38: 1283-7. [\[PubMed\]](#) [\[Google Scholar\]](#)
56. Kumar MS, Sajjan GS, Satish K, Varma KM. A comparative evaluation of efficacy of protaper universal rotary retreatment system for gutta-percha removal with or without a solvent. *Contemp Clin Dent.* 2012;3 (Suppl 2):S160-3. [\[PubMed\]](#) [\[Google Scholar\]](#)
57. B. Marques da Silva, F. Baratto-Filho, D. P. Leonardi, A. Henrique Borges, L. Volpato, F. Branco Barletta. Effectiveness of ProTaper, D-RaCe, and Mtwo retreatment files with and without supplementary instruments in the removal of root canal filling material. *Int Endod J* 2012; 45: 927-32. [\[PubMed\]](#) [\[Google Scholar\]](#)
58. Ma J, Al-Ashaw AJ, Shen Y, Gao Y, Yang Y, Zhang C, Haapasalo M. Efficacy of ProTaper Universal Rotary Retreatment system for gutta-percha removal from oval root canals: a micro-computed tomography study. *J Endod.* 2013; 38: 1516-20. [\[PubMed\]](#) [\[Google Scholar\]](#)
59. Beasley RT, Williamson AE, Justman BC, Qian F. Time required to remove guttacore, thermafil plus, and thermoplasticized gutta-percha from moderately curved root canals with protaper files. *J Endod.* 2013; 39: 125-8. [\[PubMed\]](#) [\[Google Scholar\]](#)
60. Zuolo AS, Mello JE Jr, Cunha RS, Zuolo ML, Bueno CE. Efficacy of reciprocating and

rotary techniques for removing filling material during root canal retreatment. Int Endod J. 2013 ; 46: 947-53. [\[PubMed\]](#) [\[Google Scholar\]](#)

61. Vale MS, Moreno Mdos S, Silva PM, Botelho TC. Endodontic filling removal procedure: an ex vivo comparative study between two rotary techniques. Braz Oral Res. 2013; 27: 478-83. [\[PubMed\]](#) [\[Google Scholar\]](#)
62. Mittal N, Jain J. Spiral computed tomography assessment of the efficacy of different rotary versus hand retreatment system. J Conserv Dent. 2014; 17: 8-12. [\[PubMed\]](#) [\[Google Scholar\]](#)
63. Topçuoğlu HS, Demirbuga S, Tuncay Ö, Pala K, Arslan H, Karataş E. The effects of Mtwo, R-Endo, and D-RaCe retreatment instruments on the incidence of dentinal defects during the removal of root canal filling material. J Endod. 2014; 40: 266-70. [\[PubMed\]](#) [\[Google Scholar\]](#)
64. Rached-Júnior FA, Sousa-Neto MD, Bruniera JF, Duarte MA, Silva-Sousa YT. Confocal microscopy assessment of filling material remaining on root canal walls after retreatment. Int Endod J. 2014; 47: 264-70. [\[PubMed\]](#) [\[Google Scholar\]](#)
65. Rios Mde A, Villela AM, Cunha RS, Velasco RC, De Martin AS, Kato AS, Bueno CE. Efficacy of 2 reciprocating systems compared with a rotary retreatment system for gutta-percha removal. J Endod. 2014; 40: 543-6. [\[PubMed\]](#) [\[Google Scholar\]](#)
66. Iriboz E, Sazak Öveçoğlu H. Comparison of ProTaper and Mtwo retreatment systems in the removal of resin-based root canal obturation materials during retreatment. Aust Endod J. 2014; 40: 6-11. [\[PubMed\]](#) [\[Google Scholar\]](#)
67. Topçuoğlu HS, Düzgün S, Kesim B, Tuncay O. Incidence of apical crack initiation and propagation during the removal of root canal filling material with ProTaper and Mtwo rotary nickel-titanium retreatment instruments and hand files. J Endod. 2014; 40: 1009-12. [\[PubMed\]](#) [\[Google Scholar\]](#)
68. Paik S, Sechrist C, Torabinejad M. Levels of evidence for the outcome of endodontic retreatment. J Endod. 2004; 30: 745-50. [\[PubMed\]](#) [\[Google Scholar\]](#)
69. Basrani B, Haapasalo M. Update on endodontic irrigating solutions. Endod Topic 2012; 2: 74-102. [\[Google Scholar\]](#)
70. Lee KW, Williams MC, Camps JJ, Pashley DH. Adhesion of endodontic sealers to dentin and gutta-percha. J Endod. 2002; 28: 684-8. [\[PubMed\]](#) [\[Google Scholar\]](#)
71. Mamootil K, Messer HH. Penetration of dentinal tubules by endodontic sealer cements in extracted teeth and in vivo. Int Endod J. 2007; 40: 873-81. [\[PubMed\]](#) [\[Google Scholar\]](#)
72. Lea CS, Apicella MJ, Mines P, Yancich PP, Parker MH. Comparison of the obturation density of cold lateral compaction versus warm vertical compaction using the continuous wave of condensation technique. J Endod. 2005; 31: 37-9. [\[PubMed\]](#) [\[Google Scholar\]](#)
73. Al-Ali M, Sathorn C, Parashos P. Root canal debridement efficacy of different final irrigation protocols. Int Endod J. 2012; 45: 898-906. [\[PubMed\]](#) [\[Google Scholar\]](#)
74. Rödiger T, Sedghi M, Konietzschke F, Lange K, Ziebolz D, Hülsmann M. Efficacy of syringe irrigation, RinsEndo and passive ultrasonic irrigation in removing debris from irregularities in root canals with different apical sizes. Int Endod J. 2010; 43: 581-9. [\[PubMed\]](#) [\[Google Scholar\]](#)
75. Jiang LM, Verhaagen B, Versluis M, Zangrillo C, Cuckovic D, van der Sluis LW. An evaluation of the effect of pulsed ultrasound on the cleaning efficacy of passive ultrasonic irrigation. J Endod. 2010; 36: 1887-91. [\[PubMed\]](#) [\[Google Scholar\]](#)
76. Patel S. New dimensions in endodontic imaging: Part 2. Cone beam computed tomography. Int Endod J. 2009; 42: 463-75. [\[PubMed\]](#) [\[Google Scholar\]](#)
77. Robertson D, Leeb IJ, McKee M, Brewer E. A clearing technique for the study of root canal systems. J Endod. 1980; 6: 421-4. [\[PubMed\]](#) [\[Google Scholar\]](#)



Avances en Biomedicina se distribuye bajo la Licencia Creative Commons Atribución-NoComercial-CompartirIgual 4.0 Venezuela, por lo que el envío y la publicación de artículos a la revista son completamente gratuitos.



<https://me-qr.com/Jfqxc09L>

How to cite this article.

Upendranatha Reddy N, Gopi Krishna Reddy M, Sujayeendranatha Reddy E, Anusha Y, Nagalakshmi Reddy S, Datta Sai Kiran K. Evaluation of the efficacy of Protaper, HERO SHAPER GOLD, ProTaper Universal retreatment, and R-Endo, for removing gutta-percha and AH plus sealer from root canals. *Avan Biomed* 2025; 14: XX